

Supplementary Table 1

Odds ratios of individual DSM-IV/V symptoms in predicting ADHD diagnosis at different times

<u>DSM-IV Symptom</u>	<u>A: T1</u>			<u>B: T2</u>			<u>C: T3</u>		
	<u>O.R.</u>	<u>95% C.I.</u>	<u>R.O.</u>	<u>O.R.</u>	<u>95% C.I.</u>	<u>R.O.</u>	<u>O.R.</u>	<u>95% C.I.</u>	<u>R.O.</u>
1. Fails to give close attention, careless	7.3	3.1-16.9	17	15.8	6.7-37.0	11	31.5	12.1-82.2	5
2. Difficulty sustaining attention	13.7	5.8-32.0	12	17.8	7.4-42.9	9	40.6	14.7-111.7	4
3. Does not listen	23.8	9.3-61.2	5	17.5	7.2-42.5	10	21.3	8.8-51.7	10
4. Does not follow through	12.7	5.3-30.4	13	24.7	9.6-63.4	5	40.9	15.3-109.133	3
5. Difficulty organizing	10.7	4.4-25.8	15	11.9	5.2-27.2	15	20.7	8.7-49.5	11
6. Avoids tasks	3.4	1.5-8.0	18	10.9	4.7-25.2	16	9.1	4.0-20.4	16
7. Loses things	11.4	4.5-28.9	14	13.2	5.6-30.9	14	27.7	9.7-79.2	7
8. Easily distracted	19.9	7.8-50.9	6	78.0	17.5-347.1	1	30.6	10.8-86.4	6
9. Forgetful	26.4	7.2-96.2	3	13.9	5.7-33.8	13	43.8	12.3-155.9	2
10. Fidgets or squirms	77.0	17.2-344.5	1	31.0	10.9-88.1	2	53.9	18.3-159.0	1
11. Leaves seat	27.7	9.7-78.6	2	16.7	7.0-39.6	11	16.1	6.9-37.6	14
12. Runs or climbs	24.4	9.6-61.7	4	21.1	8.7-51.6	8	19.3	7.5-49.5	12
13. Difficulty playing quietly	17.2	5.8-50.4	10	23.7	7.5-74.6	4	24.3	6.8-86.0	8
14. "On the go"	18.2	7.5-44.2	7	21.7	8.7-54.2	7	22.1	7.8-63.0	9
15. Talks excessively	9.3	4.1-21.0	16	3.4	1.7-7.1	18	7.5	3.5-16.1	18
16. Blurts out answers	17.6	7.3-42.7	8	10.0	4.5-22.2	17	10.4	4.5-24.2	15
17. Difficulty awaiting turn	17.6	7.3-42.7	9	15.8	6.6-37.5	12	16.6	6.5-42.4	13
18. Interrupts or intrudes	15.8	6.0-41.6	11	23.4	8.3-65.4	6	8.2	3.8-17.6	17

Note. T1=Time 1; T2=Time 2; T3=Time 3; O.R.=Odds Ratio; 95% C.I.=95% Confidence Interval; R.O.=Rank Order.