

Supplemental table 2: HADS Individual Item Scores for SST Participants in Whom CNV Developed in the Second Eye															
Before the last SST examination, Comparing Scores from (1) Baseline to prior to CNV diagnosis, and (2) Prior to and following CNV diagnosis															
Comparison of mean and median HADS scores															
Baseline to prior to CNV diagnosis								Prior to and following CNV diagnosis							
HADS Item	N	Baseline Mean	Baseline Median	Prior Mean	Prior Median	p-value*	Adjusted p-value**	N	Prior Mean	Prior Median	Following Mean	Following Median	p-value*	Adjusted p-value**	
hads_tense	66	3.44	4.00	3.50	4.00	0.50	1.00	77	3.43	4.00	3.55	4.00	0.17	1.00	
hads_enjoy	66	1.47	1.00	1.61	1.00	0.32	1.00	77	1.58	1.00	1.82	2.00	0.04	0.53	
hads_awful	66	3.56	4.00	3.64	4.00	0.43	1.00	77	3.62	4.00	3.66	4.00	0.60	1.00	
hads_laugh	66	1.33	1.00	1.32	1.00	0.89	1.00	77	1.36	1.00	1.30	1.00	0.53	1.00	
hads_worrying	62	3.52	4.00	3.39	3.00	0.23	1.00	66	3.30	3.00	3.59	4.00	0.006	0.08	
hads_cheerful	66	3.83	4.00	3.68	4.00	0.05	0.69	77	3.79	4.00	3.71	4.00	0.32	1.00	
hads_relaxed	66	1.67	2.00	1.56	1.00	0.20	1.00	77	1.53	1.00	1.65	2.00	0.25	1.00	
hads_slowed_down	66	2.94	3.00	2.92	3.00	0.88	1.00	77	2.94	3.00	2.86	3.00	0.50	1.00	
hads_butterflies	66	1.44	1.00	1.33	1.00	0.20	1.00	77	1.34	1.00	1.30	1.00	0.57	1.00	
hads_appearance	66	3.82	4.00	3.85	4.00	0.66	1.00	77	3.84	4.00	3.91	4.00	0.23	1.00	
hads_restless	66	3.33	3.00	3.38	4.00	0.63	1.00	77	3.39	4.00	3.29	3.00	0.30	1.00	
hads_look_forward	66	1.45	1.00	1.50	1.00	0.66	1.00	77	1.49	1.00	1.47	1.00	0.78	1.00	
hads_panic	66	3.64	4.00	3.70	4.00	0.42	1.00	77	3.66	4.00	3.64	4.00	0.73	1.00	
hads_book	66	1.15	1.00	1.33	1.00	0.08	1.00	77	1.43	1.00	1.65	1.00	0.08	1.00	
* Paired t-test															
** Bonferroni adjustment for multiple comparisons															
Bold font indicates statistical significance (p value < 0.05)															